

FORM PTO-1390  
(REV. 10-2000)U.S. DEPARTMENT OF COMMERCE  
PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

20342P

**TRANSMITTAL LETTER TO THE UNITED STATES  
DESIGNATED/ELECTED OFFICE (DO/EO/US)  
CONCERNING A FILING UNDER 35 U.S.C. 371**

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)

**09/ 857917**INTERNATIONAL APPLICATION NO.  
PCT/US99/30297INTERNATIONAL FILING DATE  
20 DEC 1999PRIORITY DATE CLAIMED  
23 DEC 1998

## TITLE OF INVENTION

NEUTRALIZING ASSAY USING HUMAN PAPILLOMAVIRUS VIRUS-LIKE PARTICLES

## APPLICANT(S) FOR DO/EO/US

KATHRIN U. JANSEN, MARK D. YEAGER, PAUL M. KELLER, ROBERT S. LOWE, MIGUEL ASTE-AMEZAGA

**Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:**

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☐ This is an express request to begin national examination procedures [35 U.S.C. 371(f)] at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. ☐ A proper Demand for International Preliminary Examination was made and the US was elected by the expiration of the 19th month from the earliest claimed priority date (PCT Article 31).
5. ☒ A copy of the International Application as filed [35 U.S.C. 371(c)(2)].
  - a. ☐ is attached hereto (required only if not communicated by the International Bureau).
  - b. ☐ has been communicated by the International Bureau.
  - c. ☒ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ An English language translation of the International Application as filed [35 U.S.C. 371(c)(2)].
- ☒ Amendments to the claims of the International Application under PCT Article 19 [35 U.S.C. 371(c)(3)].
  - a. ☐ are attached hereto (required only if not communicated by the International Bureau).
  - b. ☐ have been communicated by the International Bureau.
  - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
  - d. ☒ have not been made and will not be made.
8. ☐ An English language translation of the amendments to the claims under PCT Article 19 [35 U.S.C. 371(c)(3)].
9. ☒ An oath or declaration of the inventor(s) [35 U.S.C. 371(c)(4)].
10. ☐ An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 [35 U.S.C. 371(c)(5)].

**Items 11 to 16 below concern other document(s) or information included:**

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☒ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☐ A **FIRST** preliminary amendment.
- ☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☐ Other items or information:

**EXPRESS MAIL CERTIFICATE**  
 DATE OF DEPOSIT June 11, 2001  
 EXPRESS MAIL NO. EL520788529 US  
 I HEREBY CERTIFY THAT THIS CORRESPONDENCE IS  
 BEING DEPOSITED WITH THE UNITED STATES POSTAL  
 SERVICE AS EXPRESS MAIL. "YOUR OFFICE TO  
 ADDRESSEE" BEFORE 5 PM ON THE ABOVE DATE IN  
 AN ENVELOPE ADDRESSED TO ASSISTANT COMMISSIONER  
 FOR PATENTS, WASHINGTON, D.C. 20231.  
 MAILED BY P. B. Crowley  
 DATE 6/14/01

U.S. APPLICATION NO. (If known, see 37 CFR 1.5) <b>09/857917</b>		INTERNATIONAL APPLICATION NO. PCT/US99/30297		ATTORNEY'S DOCKET NUMBER 20342P	
---	--	---	--	------------------------------------	--

17. <input checked="" type="checkbox"/> The following fees are submitted: <b>BASIC NATIONAL FEE [37 CFR 1.492(a)(1)-(5)]:</b> Neither international preliminary examination fee (37 CFR 1.482) nor international search fee [37 CFR 1.445(a)(2)] paid to USPTO and International Search Report not prepared by the EPO or JPO..... <b>\$1,000.00</b> International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO..... <b>\$860.00</b> International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee [37 CFR 1.445(a)(2)] paid to USPTO..... <b>\$710.00</b> International preliminary examination fee paid to USPTO (37 CFR 1.482) but all claims did not satisfy provisions of PCT Article 33(1)-(4)..... <b>\$690.00</b> International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(1)-(4)..... <b>\$100.00</b> <div style="text-align: right;"><b>ENTER APPROPRIATE BASIC FEE AMOUNT =</b></div>				<b>CALCULATIONS</b>		<b>PTO USE ONLY</b>	

Surcharge of <b>\$130.00</b> for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date [37 CFR 1.492(e)].				\$0.00			
--	--	--	--	--------	--	--	--

Claims	Number Filed	Number Extra	Rate			
Total Claims	12 - 20 =	0	X <b>\$18.00</b>		\$0.00	
Independent Claims	3 - 3 =	0	X <b>\$80.00</b>		\$0.00	
Multiple dependent claim(s) (if applicable)			+ <b>\$270.00</b>		\$0.00	
<b>TOTAL OF ABOVE CALCULATIONS =</b>					\$100.00	

<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2.							
<b>SUBTOTAL =</b>					\$100.00		

Processing fee of <b>\$130.00</b> for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date [37 CFR 1.492(f)].				\$0.00			
<b>TOTAL NATIONAL FEE =</b>					\$100.00		

Fee for recording the enclosed assignment [37 CFR 1.21(h)]. The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). <b>\$40.00</b> per property.				\$0.00			
<b>TOTAL FEES ENCLOSED =</b>					\$100.00		
				Amount to be refunded			
				charged			

a. ☐ A check in the amount of \$ \_\_\_\_\_ to cover the above fees is enclosed.

b. ☒ Please charge my Deposit Account No. 13-2755 in the amount of \$100.00 to cover the above fees. A duplicate copy of this sheet is enclosed.

c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to the Deposit Account No. 13-2755. A duplicate copy of this sheet is enclosed.


**NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive [37 CFR 1.137(a) or (b)] must be filed and granted to restore the application to pending status.**

SEND ALL CORRESPONDENCE TO:

MERCK & CO., INC.  
 Patent Department, RY60-30  
 P.O. Box 2000  
 126 East Lincoln Avenue  
 Rahway, New Jersey 07065-0970

DATE: June 11, 2001

PHONE #: (732) 594-3046

  
 SIGNATURE  
 JOANNE M. GIESSER  
 NAME  
32,838  
 REGISTRATION NUMBER

NEUTRALIZING ASSAY USING HUMAN PAPILLOMAVIRUS VIRUS-LIKE  
PARTICLES

FIELD OF THE INVENTION

This invention is related to a novel method of detecting human papillomavirus (HPV) neutralizing antibodies in a serum or plasma.

BACKGROUND OF THE INVENTION

Human Papillomaviruses (HPV) infect the genital tract and has been associated with various dysplasias, cancers and other diseases. These diseases are currently targets for vaccine developments, and vaccines based on virus-like particles (VLPs) which contain only L1 or L1+L2 capsid proteins are currently in trials.

To test vaccine candidates, neutralization assays would be desirable. While there are over 100 HPV types described today, there are only a few HPV types for which neutralization assays exist. Since HPV does not grow in tissue culture, the development of neutralization assays for HPV has been difficult.

To facilitate vaccine development, it would be desirable to develop a basic neutralization assay which could be easily modified so as to have a neutralization assay for any HPV type.

DESCRIPTION OF THE INVENTION

This invention relates to a papillomavirus "pseudovirion" comprising a virus-like particle (VLP) which is covalently linked to a reporter gene DNA construct. In preferred embodiments, the VLPs are human papillomavirus (HPV) VLPs.

This invention also includes assays to determine if an anti-VLP antibody is present in a sample comprising the steps of: combining a cell culture susceptible to infection by the pseudovirions, the sample, and pseudovirions; and measuring the amount of uptake of the pseudovirions by the cell. The uptake may be compared to a control where no sample is present, or the sample is a pre-immune sample, wherein a decrease in cell uptake of pseudovirions indicates the presence of neutralizing antibodies in the sample.

Another aspect of this invention is a method of making the pseudovirion construct, comprising generating relatively pure VLPs and attaching the reporter gene construct with a heterobifunctional cross-linker.

Yet another aspect of this invention is a method of delivery of a reporter genes into a cell comprising infecting a cell with a pseudovirion.

#### DESCRIPTION OF THE DRAWINGS:

Figure 1 is a diagram showing a reporter gene construct based on the beta-lactamase gene (BLAM).

Figure 2 shows the inhibition of HPV16 pseudovirion uptake into cells by HPV16 neutralizing monoclonal antibody H16.V5.

Figure 3 shows the inhibition of HPV16 pseudovirion uptake into cells by neutralizing antibody in the sera of individuals immunized with HPV16 VLPs.

As used throughout the specification and claims, the following definitions apply:

"Pseudovirion" means a papillomavirus virus-like particle which is chemically linked or bonded to a reporter gene construct, either directly or through a linking group.

"Reporter gene construct" means any nucleic acid encoding a protein whose transcription, translation, or post-translation activity can be conveniently detected, along with known necessary elements required for control of its transcription and translation (including, for example, promoters, enhancers, transcription termination sequences, and the like). Examples of reporter gene constructs include genes for beta-galactosidase, luciferase, and green-fluorescent protein, and the like.

#### DETAILED DESCRIPTION OF THE INVENTION

The pseudovirions and methods of using them do not depend on any particular papillomavirus strain or any other viral capsid. In preferred embodiments, the papillomavirus is a human papillomavirus (HPV), and it is particularly preferred that the HPV is one of the strains which is associated with genital warts and/or genital cancers, and may be selected from the group consisting of: HPV6a, HPV6b, HPV11, HPV16, HPV18, HPV31, HPV33, HPV35, HPV42, HPV43, HPV44, HPV45, HPV51, HPV52, and HPV56.

A pseudovirion can be made from virtually any papillomavirus VLP. A wild-type VLP generally is made up predominantly of L1 and a minor amount of L2 protein. However, it is known that VLPs may contain only L1 protein. In addition, VLPs

have been described which contain modified L1 protein, modified L2 proteins, proteins (whether or not naturally occurring, such as HPV E-proteins) which have been fused to at least a portion of the L1 or L2 proteins, and/or an additional, non-naturally occurring protein. For purposes of this invention, any of the foregoing may be used as a starting material for the construction of pseudovirions. Generally, preferred VLPs contain either L1 or the combination of L1 and L2 ("L1+L2") protein.

The L1 or L1+L2 protein-containing VLPs can be made by transforming a selected host cell with genes encoding the L1 or L1+L2 proteins. General techniques, the genes for various types of L1 and L2, and methods for recombinant expression of L1 and L2 proteins are known in the art, and may be used. Preferred host cells include yeast, insect cells, mammalian cells and *E. coli*.

Under one preferred embodiment of this invention, yeast are transformed with plasmid DNA containing the genes encoding L1 protein under control of known yeast promoters, such as the yeast Gal 1 and Gal 10 promoters. Expression of the gene product is induced by addition of galactose to the growth media, and the VLPs are isolated from the induced cell lysates.

Virtually any reporter gene construct can be coupled to the VLPs to make a pseudovirion. One preferred reporter gene construct used is the beta-lactamase gene ("BLAM"), but other reporter constructs encoding enzymes or proteins like beta-galactosidase, luciferase and green-fluorescent protein or genes coding for spliced transcripts can also be used.

The reporter construct may be coupled to the VLP using known coupling reactions. In a preferred embodiment, the reporter gene construct is attached to a linker which is commercially available, and the linker is attached to the VLP. A preferred cross-linking reagent is (succinimidyl-maleimidomethyl-cyclohexane-carboxylate "SMCC"). Other potentially useful crosslinking reagents in addition to SMCC would have the following properties: 1) heterobifunctional or sequentially activated crosslinkers that would eliminate interfering oligo-oligo or VLP-VLP crosslinking, 2) stable reactions with either 5'-amino or 5'-thiol modified oligonucleotides, 3) stable reactions with primary amine, free sulfhydryl, or carbohydrate moieties on the VLP protein. These reagents would include, but are not limited to the following and their sulfated derivatives: SMPB (succinimidyl-4-(p-maleimidophenyl) butyrate), SIAB (succinimidyl(4-iodoacetyl) amino benzoate), GMBS (n-(4-maleimido-butyryloxy)-succinimide), MBS (m-maleimidobenzoyl sulfosuccinimide ester), and MPBH (4-(4-N-maleimidophenyl) butyric acid hydrazide hydrochloride).

In constructing such a linker, molar ratio of reporter gene constructs to the VLP may vary. In accordance with this invention, the ratio of reporter gene construct to VLP should be from about 0.1 to 50, preferably from about 1 to 5, and more preferably substantially about 1 to 1.

One of the advantages of this invention is that pseudovirions can be produced easily for a variety of HPVs using the same reporter and test systems. Another advantage is that the reporter systems can be easily modified, if desired.

An important aspect of this invention is the use of pseudovirions to detect the presence of neutralizing antibodies. Although some information is known about the biology of HPV and other papillomaviruses, they have not yet been successfully cultured *in vitro*. Further, while a large number of animal species are susceptible to papillomavirus infection, each species is only susceptible to its own strains of papillomavirus. Thus, for example, while cottontail rabbit papillomavirus (CRPV) can be successfully studied in the cottontail rabbit and used as a model for the human disease, human papillomavirus cannot be directly studied in the rabbit. Various vaccines based on HPV VLPs are currently under development, but to date neutralization assays have not been available for all strains of HPV.

In the assay of this invention, a pseudovirion, a sample suspected of containing neutralizing antibodies, and a cell culture are brought into contact. The cells used in this assay may be any cells known to be susceptible to papillomavirus infection. For HPV pseudovirions, the cells may be human carcinoma cells, such as C33A (available from the ATCC), HeLa cells, and the like. If the sample (which in preferred embodiments may be serum from a patient), contains neutralizing antibodies, then the neutralizing antibodies will interact with the pseudovirions and will interfere with the uptake of the pseudovirions the cell. Optionally, a control assay may be performed, wherein the same pseudovirions and cells are used, but the sample is either eliminated, or a sample which is known not to contain neutralizing antibodies is substituted. Another example of a control sample is one from a preimmune source.

In the assay, the decreased uptake of pseudovirions into the cell is correlated to the amount of neutralizing antibody present in the sample. Depending on the reporter gene construct employed, the amount may be quantified. Quantitation of the number of cells expressing the reporter gene, and the extent of expression, can be realized using flow cytometry. Simultaneous measurement of the 520nm/450nm ratio of emission wavelengths is directly related to the proportion of cleaved CCF2 substrate in individual cells, and is

independent of substrate concentration or cell dimensions, thus resulting in better signal to noise than is obtainable with either wavelength alone.

In addition, subpopulations of cells that respond to infection could be enriched through fluorescence-activated cell sorting and cultured to expand the population of responsive cells. for example using a quantitative RT PCR assay of a reporter splice variant. These assays of this invention may be miniaturized and/or automated, and used in a 96-well format for high-throughput screens.

The invention also allows labeling of pseudovirions with different reporters to test multivalent neutralizing sera on a mixture of these pseudovirions. This can be used to follow a multivalent vaccine response in a subject.

Another aspect of this invention is the use of pseudovirions to carry different genetic materials into a variety of target cells. Substantially the same linker system is used to construct the pseudovirion, but the reporter gene construct is replaced with a gene construct which expresses a gene of choice.

The following non-limiting Examples are presented to illustrate the invention better.

### EXAMPLE 1

#### Expression of HPV16 VLPs in yeast

HPV 16 VLPs are made essentially as described in published PCT application WO95/31532, which is hereby incorporated by reference.

#### Cloning of HPV16 L1 and L2 Genes

Total genomic DNA was extracted from Caski cells (ATCC #CRL 1550) by standard techniques. The DNA was digested with Bst1107I and SphI endonucleases and electrophoresed through a 0.8% low-melting-temperature agarose preparative gel. A region corresponding to DNA of approximately 3.5-kbp was excised from the gel and the agarose was digested with Gelase<sup>TM</sup> enzyme (Epicentre Technologies, Inc.). The gel-purified DNA was made blunt-ended with T4 DNA polymerase and ligated with blunt-ended, phosphorylated oligodeoxynucleotide linkers that contain a buried HindIII site. The ligation mixture was digested to completion with HindIII and the approximately 3.5-kbp DNA was size-fractionated through an agarose gel as described above. The gel-purified

DNA was ligated with pUC18 plasmid DNA that had been digested with HindIII and dephosphorylated. Following transformation of competent *E. coli* DH5 cells (BRL), the plasmid library was screened for HPV16-positive clones by colony hybridization using an antisense <sup>32</sup>P-labeled oligodeoxynucleotide that is complementary to the 3'-end of the HPV-16 L1 gene (5'-GAG AGA TCT TAC AGC TTA CGT TTT TTG CGT TTA GC-3') (SEQ. ID. NO:1). A pUC18 plasmid containing a 3.3-kbp HPV16 genomic fragment was isolated and characterized by restriction enzyme and Southern blot analyses. This plasmid was designated pUC18-HPV16 L1/L2 and contains all of the L1 and L2 coding DNA sequences. Plasmid DNA was prepared using the Qiagen<sup>TM</sup> Plasmid Maxi kit (Qiagen, Inc.).

#### Construction of HPV16 L1 Yeast Expression Vector

The clone, pUC18-HPV16 L1/L2 was used as a template for PCR. The HPV16 L1 gene was amplified by PCR using Vent polymerase (New England Biolabs, Inc.), 10 cycles of amplification (94°C, 1 min; 48°C, 1 min; 72°C, 1 min 45 sec), and the following oligodeoxynucleotide primers which contain flanking BglII sites (underlined):

sense primer: 5'-CTC AGA TCT CAC AAA ACA AAA TGT CTC TTT GGC TGC CTA TGT AGG CC-3'(SEQ ID NO: 2)

antisense primer: 5'-GAG AGA TCT TAC AGC TTA CGT TTT TTG CGT TTA GC-3' (SEQ ID NO:3)

The sense primer introduces a yeast non-translated leader sequence immediately upstream to the HPV16 L1 initiating methionine codon (highlighted in bold print). The 1.5-kbp L1 PCR product was digested with BglII, gel-purified, and ligated with the BamHI digested pC1/1-GAL vector. A pC1/1-GAL plasmid was isolated containing the HPV16 L1 gene and designated, p14049-37-1. The L1 gene in p14049-37-1 was sequenced using the PRISM<sup>TM</sup> kit (ABI, Inc.) and an ABI Sequencer Model #373A according to the manufacturer's directions. The L1 gene in this isolate was shown to contain 3 nucleotide changes from the corrected, published prototype sequence (Kirnbauer, R. et al. (1993) *J. Virol.* 67: 6929-6936), resulting in two amino acid changes: His-202 to Asp; Thr-266 to Ala. Sequence analysis of the original template DNA confirmed that these changes were also present in the genomic clone pUC18-HPV16 L1/L2 and were not introduced by the PCR.



### Construction of the HPV16 L1 and L2 Yeast Expression Vector

Plasmid p14049-37-1 (pC1/1-GAL+HPV16 L1) was digested with *Sma*I, which cuts between the *GAL10* promoter and the *ADHI* transcription terminator. The 1.4-kbp HPV16 L2 gene was amplified by PCR using the pUC18-HPV16 L1/L2 DNA as template, Vent polymerase (New England Biolabs, Inc.), 10 cycles of PCR amplification (94°C, 1 min; 48°C, 1 min; 72°C, 1 min 45 sec) and the following oligodeoxynucleotide primers which contain flanking *Sma*I sites (underlined):

sense primer: 5'-TCC CCC GGG CAC AAA ACA AAA TGC GAC ACA AAC GTT  
CTG CAA AAC-3'(SEQ ID NO:4)

antisense primer: 5'-TCC CCC GGG CTA GGC AGC CAA AGA GAC ATC TGA-  
3'(SEQ ID NO:5)

The sense primer introduces a yeast non-translated leader sequence immediately upstream to the HPV16 L2 initiating methionine codon (highlighted in bold print). The 1.4-kbp L2 PCR product was digested with *Sma*I, gel-purified, and ligated with the *Sma*I digested p14049-37-1 vector. A pC1/1-GAL plasmid containing both the HPV16 L1 and L2 genes was isolated and designated, p14049-42-2. The L2 gene in p14049-42-2 was sequenced using the PRISM™ kit (ABI, Inc.) and an ABI Sequencer Model #373A according to the manufacturer's directions. The L2 gene in this isolate was shown to contain 5 nucleotide changes from the corrected, published prototype sequence (Kimbauer, R. et al. (1993) *supra*), resulting in one amino acid change: Ser-269 to Pro. Sequence analysis of the genomic clone pUC18-HPV16 L1/L2 confirmed that this change was also present in the original template DNA and was not introduced by the PCR.

### Expression of HPV16 L1 and Co-Expression of HPV16 L1 and L2 in Yeast

Plasmids p14049-37-1 (pC1/1-GAL+HPV16 L1) and p14049-42-2 (pC1/1-GAL+HPV16 L1 and L2) were used to transform *S. cerevisiae* strain #1558. The resulting recombinant strains were #1678 (HPV16-L1) and #1679 (HPV16 L1+L2) as shown in the table. Clonal isolates were grown at 30°C in YEHD medium containing 2% galactose for 68-78 hours. After harvesting the cells, the cell pellets were broken with glass beads and cell lysates analyzed for the expression of HPV16 L1 or HPV16 L2 protein by immunoblot analysis. Samples containing 40 mcg of total cellular protein were electrophoresed on 10% Tris-Glycine gels under reducing and denaturing conditions and electroblotted onto

nitrocellulose filters. The HPV16 L1 protein was immunodetected using rabbit polyclonal antisera raised against a trpE-HPV11 L1 fusion protein (D. Brown et al., *Virology* 201:46-54) as primary antibody and HRP-linked, donkey anti-rabbit IgG antibody (Amersham, Inc.) as the secondary antibody. The filters were processed using the chemiluminescent ECL Detection kit (Amersham, Inc.). A 50-55 kDa L1 protein band was detected in all samples except the negative control (pC1/1 without L1 or L2 gene). The L2 protein was detected as a 70 kDa protein band by immunoblot using mouse anti-HPV16 L2 sera raised against trpE-L2 fusion proteins expressed in *E. coli* as primary antibody. Goat anti-mouse IgG HRP-linked (Amersham, Inc.) was used as secondary antibody and the filters were processed as described above.

## EXAMPLE 2

### Purification of VLPs

Purification was done essentially as described in co-pending U.S. Provisional Patent Application 60/096,568, which is hereby incorporated by reference.

Yeast cells transformed to express VLPs were harvested and frozen for storage at -70°C. Frozen yeast cell suspension was removed from storage and thawed for approximately 3 hours at room temperature followed by approximately 18 hours at 4°C. BENZONASE® (Nycomed Pharma A/S, Copenhagen, Denmark) ( $2.8 \times 10^5$  Units/mL and 0.21 mg protein/mL) was added to the cell suspension to a final concentration of 750 Units per gram of wet cell weight, and in one experiment was reduced to 335 Units per gram wet cell weight. Cells were stirred for 15 minutes, then disrupted by two passes through a sanitized APV Gaulin 30CD homogenizer at chamber pressures of 14,500 to 16,000 psi, resulting in 95% cell disruption. The remaining lysate was gently stirred for 18 hours at 4°C.

Clarification by microfiltration. Cell lysate was clarified by cross-flow microfiltration in a diafiltration mode as follows. Lysate was transferred to a sterile process tank with a 1-inch diameter inlet and outlet ports. The microfilter was a 0.65 micron pore size hollow-fiber filter cartridge of 5 square feet surface area (A/G Technologies #CFP-6-D-8A, Needham, MA) housed in an A/G Technologies FlexStand® Benchtop Pilot Hollow Fiber system. The retentate was diafiltered with 3 volumes of Diafiltration Buffer (below) to produce the clarified lysate. Diafiltration Buffer was 0.2M (Na<sup>+</sup>) MOPS, pH 7.0 + 0.4M NaCl.

Chromatography of clarified lysate. The clarified lysate was fractionated by column chromatography using POROS® 50HS strong cation-exchange chromatography resin (PerSeptive Biosystems, Framingham, MA) packed in a chromatography column. The column was sanitized with 0.5 N NaOH prior to use. The column was equilibrated with HPV Diafiltration Buffer [0.2M (Na<sup>+</sup>)MOPS, pH 7.0 + 0.4M NaCl] at room temperature. The cold (4°C) clarified lysate was pumped onto the column at 125 mL/minute and the column was washed with 8 column volumes of room temperature HPV Column Buffer A [0.05M (Na<sup>+</sup>)MOPS, pH 7.0 + 0.5M NaCl] at 125 mL/minute with a linear gradient of 100% HPV Column Buffer A to 100% HPV Column Buffer B [0.05M (Na<sup>+</sup>)MOPS, pH 7.0 + 1.5 M NaCl]. Total linear gradient was 10 column volumes and was collected in 10 equal-volume fractions. Following the gradient, the column was washed with two column volumes of room temperature HPV Column Buffer B at 125 mL/minute which were collected in two additional fractions. Fractions were collected in sterile 2-liter plastic bottles and stored at 4°C. Fractions containing the last UV-absorbing peak (A280nm and A230 nm) in the gradient were pooled, filtered using a MILLIPAK-200 disposable filter unit (Millipore, Bedford, MA) and stored at 4°C.

Hydroxyapatite Chromatography All steps were carried out at room temperature. A chromatography column (13 mm ID x 36 mm) packed with Ceramic Hydroxyapatite, Type II (BioRad Cat.#7320081, Hercules, CA), was pre-equilibrated in 50 mM MOPS, pH 7.0 + 1.25 M NaCl. The partially purified HPV solution from the previous step was applied to the column at a linear flow rate of 90 cm/hour. After sample application was complete, the column was washed with eight column volumes of pre-equilibration buffer until the optical density of the column effluent was nearly zero. The HPV product was eluted with a 0% to 100% linear gradient of elution buffer (0.2 M sodium phosphate, pH 7.0 + 1.25M NaCl), also at a linear flow rate of 90 cm/hour. The total volume of the gradient was four column volumes. Fractions containing the vaccine product were identified by RIA and Bradford protein assay. The protein concentration of the product was 100 µg/mL.

Assays : Bradford protein assays were performed using Coomassie Plus Assay Reagent (Pierce, Rockford, IL) using bovine serum albumin (BSA) as a standard. Lowry protein assays were performed according to the procedure of Lowry et al 1951 *J. Biol. Chem.* 193:265-270 using BSA as a calibration standard. Antigen was assayed by a multilayered ELISA using a monoclonal antibody that recognized a conformational epitope of the VLP. Microtiter plates were coated with polyclonal goat anti-HPV VLP antibodies.

Standard and test samples were diluted with PBS containing 1% w/v BSA, 0.1% TWEEN-20, and 0.1% sodium azide and were added to the wells where antigen was captured by the plate-bound antibodies. Monoclonal anti-HPV L1 VLP antibody (Chemicon, Temecula, CA) was added to the wells to bind the antigen captured by the plate-bound antibodies. The monoclonal anti-HPV VLP antibodies were detected by horseradish peroxidase-conjugated anti-mouse IgG antibodies. A chromogenic substrate for horseradish peroxidase, 3,3',5,5'-tetramethylbenzidine (Pierce) was added and absorbance at 450 nm was proportional to the concentration of L1 VLP in the sample.

### EXAMPLE 3

#### Generation of the beta-lactamase reporter and coupling to HPV16 VLPs to produce "pseudovirions"

The beta-lactamase reporter gene was constructed as follows:  
5'-amine modified oligos were designed, synthesized (Midland Certified Reagents; Midland, TX) and used to amplify the section of the Aurora vector pcDNA3-BLAM (Zlokarnik et al 1998 *Science* 279:84-88) containing the CMV promoter + BLAM + poly-A tail (Figure 1). Primers for both coupling orientations can be used (i.e., amine at 5' end of sense strand or amine at 5' end of antisense strand). The downstream primers were 5'-biotinylated in order to reduce degradation by DNA exonuclease in the final assay, and to provide the ability to detect DNA. For primer sequences see Table 1.

Table 1.

Primer Name	Primer Sequence
pcdna3.seq-182F.5AM (A)	5'-amino-TAGGGTTAGG-CGTTTTGC-3' (SEQ.ID.NO:6)
pcdna3.seq-182F.5bio (B)	5'-biotin-TAGGGTTAGG-CGTTTTGC-3' (SEQ.ID.NO:7)
pcdna3.seq-1266R.5AM (C)	5'-amino-CCATAGAGCC CACCGC (SEQ.ID.NO:8)
pcdna3.seq-1266R.5bio (D)	5'-biotin-CCATAGAGCC CACCGC (SEQ.ID.NO:9)

Primers A and B were initially used to amplify a 1798 base pair (bp) sequence encoding the 591 bp human CMV promoter, the 809 bp beta-lactamase and the 135 bp bovine growth hormone poly A terminator sequences. PCR products were purified using gel filtration (spin-columns with a silica-gel membrane, QIAGEN, Inc.), quantitated by UV spectrometry at OD 260 nm, and conjugated to the NHS-ester of the heterobifunctional crosslinker SMCC (succinimidyl-maleimidomethyl-cyclohexane-carboxylate) (Pierce) according to the manufacturer's instructions for 1 hr at 20°C.

The conjugated reporter gene was purified by gel filtration (Microspin G-25, Pharmacia), then reacted to free sulfhydryls on concentrated HPV 16 VLPs (0.97 mg/mL), via the maleimide group (approx. 1 free sulfhydryl per L1 monomer).

The pseudovirions were then sterile filtered to be used in the neutralization assay. Initial experiments indicated that this low coupling ratio of reporter to VLP was beneficial (data not shown). The reaction took place at 4°C for 12 hours.

#### EXAMPLE 4

##### Infection of C33A cells with HPV pseudovirions

2000 C33 A cells (ATCC #HTB 31) were cultured for 3 days at 37°C in complete DMEM (Dulbecco's Modified Eagle Medium (GIBCO BRL) containing Penicillin and Streptomycin (100 U/mL and 100 mcg/mL, respectively) and 10% fetal calf serum. Before infection, the serum containing medium was removed and 200 ng HPV 16 pseudovirions in 50 mcL of opti-MEM (GIBCO BRL) were added. 24 hours after infection, the cells were fed with 100 mcL of complete DMEM medium. Two days after infection, beta-lactamase reporter gene expression was detected in infected CA33A cells using CCF2 (Aurora), a substrate for the beta-lactamase. (Zlokarnik, *supra*).

Briefly, cells were washed with opti-MEM medium before 50 mcL of loading buffer containing CCF2 was added to each well of a 96 well plate (FALCON, Becton-Dickinson). CCF2 final concentration was 5 microM. Plates were incubated for 3h at room temperature. Conversion of CCF2 in cells expressing beta-lactamase was followed by monitoring the shift in fluorescence emission wavelength to blue at 460 nm using a fluorescent microscope (Olympus, Mod. IX70). On average, 204 +/- 55 (n=12) blue cells could be detected two days after infection.

### EXAMPLE 5

#### Neutralization of HPV16 pseudovirion infection by monoclonal antibodies and immune sera from HPV16 VLP immunized individuals

2000 C33A cells/well (ATCC #HTB 31) were cultured in 96 well plates (FALCON, Becton-Dickinson) for 3 days at 37°C in complete DMEM (GIBCO BRL). 25 mcl of 1:40 dilutions of either (i) human pre-immune sera, (ii) immune sera from individuals immunized with HPV16 VLPs, (iii) immune sera from individuals immunized with other HPV VLPs, or (iv) various dilutions of HPV VLP-specific monoclonal antibodies were added to the wells, followed by 25 mcl of medium containing 200 ng of pseudovirions. The plates were incubated at 37°C in a CO<sub>2</sub> incubator (5% CO<sub>2</sub>) and then analyzed as described in Example 3. Percent neutralization was determined by calculating the percentage of reduction of blue cells in wells containing HPV16 neutralizing monoclonal antibodies or human immune sera versus wells containing non-specific monoclonal antibodies or pre-immune sera. Figure 2 shows the percent neutralization of HPV16 pseudovirions by serial dilutions of ascites containing the known HPV16 neutralizing monoclonal antibody H16. V5. A 1:10,000 dilution of ascites containing the nonspecific monoclonal antibody H18.J4 was not neutralizing. Using the described assay, a panel of 52 sera of individuals immunized with HPV16 VLPs was tested at a 1:80 final serum dilution. Most of the immune sera were neutralizing in this assay (Figure 3). In addition, a good correlation was seen between HPV16 VLP-specific antibody titers determined by an HPV16 RIA antibody test and % neutralization (Figure 3).

### EXAMPLE 6

Polystyrene beads (1/4 inch with specular finish) were pre-coated with normal goat serum at a 1:1000 dilution in phosphate-buffered saline (PBS). Batches of 3000 beads, 7.5 beads/ml, were incubated overnight at 4°C. Beads were then washed in 5 volumes of deionized water, blotted dry on paper towels and permitted to air dry on a stainless steel tray. HPV 16 L1 VLP were coated onto the pre-coated beads using MOPS buffer (50 mM MOPS + 0.5 M NaCl, pH 7.0) containing 50 ng of VLP per ml (5 beads per ml). Antigen and buffer were mixed first in a Corning® media bottle then the beads were added. The bottle containing the coating antigen and beads was secured onto a Cole Palmer rotator,

coating was for 1 hour at room temperature at 5 rpm. Beads were washed in 5 volumes of MOPS buffer and sufficient buffer was added to keep beads submerged. They were then stored until needed at 4°C for a maximum of 4 months.

The HPV16 L1 VLP antibody RIA is a competitive antibody assay, where antibodies compete for limited number of epitopes with a limited amount of the conformational specific monoclonal antibody H16.V5. Ascites containing H16.V5 were used at a 1:800,000 dilution in PBS + 1% bovine serum albumin + 0.05% Tween 20. Equal volumes (100mcL) of sample serum and diluted H16.V5 were mixed in a well of a 20 well Abbott assay plate. Then a single HPV16 antigen-coated bead was added to the assay well. Samples were assayed in duplicates.

Assay plates were sealed with adhesive sheets and left to incubate at room temperature 17 to 24 hours. Plates were washed with deionized water using an Abbott plate washer. The amount of mouse MAB that successfully bound to the HPV16 VLP-coated bead was detected with  $^{125}\text{I}$  [Goat anti-mouse IgG] (NEN Life Sciences, Boston, MA). Plates were sealed with fresh adhesive sheet and incubated at 37°C for 2 to 2.5 hours. Plates were again washed with deionized water. The beads were then transferred to carrier tubes and read on a gamma counter. Relative inhibition of the H16.V5 binding was compared to a standard curve using a four parameter logistic curve fit. The standard curve was derived using an HPV16 immune serum with an arbitrarily assigned number of units (mMU/mL).

## WHAT IS CLAIMED IS:

1. A human papillomavirus (HPV) pseudovirion comprising a virus-like particle (VLP) covalently linked to a reporter gene construct.
2. A pseudovirion according to Claim 1 wherein the HPV is selected from the group consisting of: HPV6a, HPV6b, HPV11, HPV16, HPV18, HPV31, HPV33, HPV35, HPV42, HPV43, HPV44, HPV45, HPV51, HPV52 and HPV56.
3. A pseudovirion according to Claim 1 wherein the VLP is comprised of L1 or L1+L2 protein.
4. A pseudovirion according to Claim 3 wherein the reporter gene construct is selected from the group consisting of: beta-lactamase gene constructs, beta galactosidase gene constructs, luciferase gene constructs and green fluorescent protein gene constructs.
5. A pseudovirion comprising a HPV VLP, a heterobifunctional crosslinker and a reporter gene construct.
6. A pseudovirion according to Claim 5 wherein the molar ratio of the reporter gene construct to VLP is from about 0.1 to 50.
7. A pseudovirion according to Claim 5, wherein the molar ratio of the reporter gene construct to VLP is about 1 to 5.
8. A pseudovirion according to Claim 5 wherein the molar ratio of the reporter gene construct to VLP is about 1 to 1.
9. An assay to determine if anti-HPV neutralizing antibodies are present in a sample, comprising the steps of:
  - a) contacting pseudovirions, a cell culture susceptible of being infected by pseudovirions, and the sample; and
  - b) determining if cell uptake of pseudovirions is inhibited, wherein inhibition of uptake is indicative of anti-HPV neutralizing antibodies in the sample.



10. An assay according to Claim 9 wherein the sample comprises serum from a human.
11. An assay according to Claim 10 further comprising the step of comparing the results to those obtained using a control assay.
12. An assay according to Claim 9 wherein the pseudovirions comprise a beta-lactamase gene construct.

1/4

Forward Primer

TAGGGTTAGG CGTTTGGCG TGCTTCGCGA TGTACGGGCC AGATATACGC GTTGACATTG 60  
ATTATTGACT AGTTATTAAAT AGTAATCAAT TACGGGGTCA TTAGTTCATA GCCCATATAT 120  
GGAGTTCCGC GTTACATAAC TTACGGTAA TGGCCCGCCT GGCTGACCGC CCAACGACCC 180  
CCGCCCATTG ACGTCAATAA TGACGTATGT TCCCATAGTA ACGCCAATAG GGACTTTCCA 240  
TTGACGTCAA TGGGTGGACT ATTTACGGTA AACTGCCCCAC TTGGCAGTAC ATCAAGTGTA 300  
TCATATGCCA AGTACGCCCC CTATTGACGT CAATGACGGT AAATGGCCCC CCTGGCATTA 360  
TGCCCCAGTAC ATGACCTTAT GGGACTTTCC TACTTGGCAG TACATCTAGG TATTAGTCAT 420  
CGCTATTACC ATGGTGATGC GGTTTTGGCA GTACATCAAT GGGCGTGGAT AGCGGTTTGA 480  
CTCACGGGGA TTTCCAAGTC TCCACCCCAT TGACGTCAAT GGGAGTTTGT TTTGGCACCA 540  
AAATCAACGG GACTTTCCAA AATGTCGTAA CAACTCCGCC CCATTGACGC AAATGGGCGG 600  
TAGGCGTGTA CGGTGGGAGG TCTATATAAG CAGAGCTCTC TGGCTAACTA GAGAACCCAC 660  
TGCTTACTGG CTTATCGAAA TTAATACGAC TCACTATAGG GAGACCCAAG CTTGGTACCA 720  
TGGACCCAGA AACGCTGGTG AAAGTAAAG ATGCTGAAGA TCAGTTGGGT GCACGAGTGG 780  
GTTACATCGA ACTGGATCTC AACAGCGGTA AGATCCTTGA GAGTTTTCGC CCCGAAGAAC 840  
GTTTTCCAAT GATGAGCACT TTAAAGTTC TGCTATGTGG CGCGGTATTA TCCCGTATTG 900

FIG.1A

2/4

ACGCCGGCA AGAGCAACTC GGTGCGCGCA TACACTATTC TCAGAATGAC TTGGTTGAGT 960  
ACTCACCAGT CACAGAAAAG CATCTTACGG ATGGCATGAC AGTAAGAGAA TTATGCAGTG 1020  
CTGCCATAAC CATGAGTGAT AACACTGCCG CCAACTTACT TCTGACAACG ATCGGAGGAC 1080  
CGAAGGAGCT AACCGCTTTT TTGCACAACA TGGGGGATCA TGTAACCTCGC CTTGATCGTT 1140  
GGGAACCGGA GCTGAATGAA GCCATACCAA ACGACGAGCG TGACACACAG ATGCCCTGTAG 1200  
CAATGGCAAC AACGTTGCGC AACTATTAA CTGGCGAACT ACTTACTCTA GCTTCCCGGC 1260  
AACAAATTAAT AGACTGGATG GAGCGGATA AAGTTCAGG ACCACTTCTG CGCTCGGCCC 1320  
TTCCGGCTGG CTGGTTTATT GCTGATAAAT CTGGAGCCGG TGAGCGTGGG TCTCGCGGTA 1380  
TCATTGCAGC ACTGGGGCCA GATGGTAAGC CCTCCCGTAT CGTAGTTATC TACACGACGG 1440  
GGAGTCAGC AACTATGGAT GAACGAAATA GACAGATCGC TGAGATAGGT GCCTCACTGA 1500  
TTAAGCATTG GTAATCTAGA GGGCCCTATT CTATAGTGTC ACCTAAATGC TAGAGCTCGC 1560  
TGATCAGCCT CGACTGTGCC TTCTAGTGC CAGCCATCTG TTGTTTGCCC CTCCCCCGTG 1620  
CCTTCCTTGA CCCTGGAAGG TGCCACTCCC ACTGTCTCTT CCTAATAAAA TGAGGAAATT 1680  
GCATCGCATT GTCTGAGTAG GTGTCAATTCT ATTCTGGGG GTGGGGTGGG GCAGGACAGC 1740  
AAGGGGGAGG ATTGGGAAGA CAATAGCAGG CATGCTGGG ATGCGGTGGG CTCCTATGG 1798

Reverse Primer

FIG.1B

3/4




FIG.2

4/4




FIG.3

**DECLARATION AND  
POWER OF ATTORNEY  
FOR UTILITY OR DESIGN  
PATENT APPLICATION  
(37 CFR 1.63)**☐ Declaration  
Submitted  
with Initial  
Filing

OR

☒ Declaration  
Submitted after Initial  
Filing (surcharge  
(37 CFR 1.16 (e))  
required)

Attorney Docket Number

20342 P

First Named Inventor

Kathrin U. Jansen et al.

**COMPLETE IF KNOWN**

Application Number

TO BE ASSIGNED

Filing Date

JUNE 11, 2001

Group Art Unit

Examiner Name

**As a below named inventor, I hereby declare that:**

My residence, post office address, and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

NEUTRALIZING ASSAY USING HUMAN PAPILLOMAVIRUS VIRUS-LIKE PARTICLES

(Title of the Invention)

the specification of which

☐ is attached hereto

OR

☒ was filed on (MM/DD/YYYY) 12/23/1998 as United States Application Number or PCT International

Application Number 60/113,388 and was amended on (MM/DD/YYYY) (if applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment specifically referred to above.

I acknowledge the duty to disclose to the Patent and Trademark Office all information known to me to be material to patentability as defined in 37 CFR 1.56.

I hereby claim foreign priority benefits under 35 U.S.C. 119(a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application Number(s)	Country	Foreign Filing Date (MM/DD/YYYY)	Attorney Docket Number	Priority Claimed?	
				YES	NO
				<input type="checkbox"/>	<input type="checkbox"/>
				<input type="checkbox"/>	<input type="checkbox"/>
				<input type="checkbox"/>	<input type="checkbox"/>
				<input type="checkbox"/>	<input type="checkbox"/>

☐ Additional foreign application numbers are listed on a supplemental priority data sheet PTO/SB/02B attached hereto:

I hereby claim the benefit under 35 U.S.C. 119(e) of any United States provisional application(s) listed below.

Application Number(s)	Filing Date (MM/DD/YYYY)	Attorney Docket Number
60/113,388	12/23/1998	20342PV

**DECLARATION AND POWER OF ATTORNEY for Utility or Design Patent Application**

I hereby claim the benefit under 35 U.S.C 120 of any United States application(s), or 365(c) of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT international application in the manner provided by the first paragraph of 35 U.S.C. 112, I acknowledge the duty to disclose information known to me to be material to patentability as defined in 37 CFR 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

U.S. Parent Application or PCT Parent Application Number	Parent Filing Date (MM/DD/YYYY)	Parent Patent Number (if applicable)

☐ Additional U.S. or PCT international application numbers are listed on a supplemental priority data sheet PTO/SB/02B attached hereto.

As a named inventor, I hereby appoint, respectively and individually, as my attorneys or agents with full power of substitution and revocation, the following registered practitioner(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

☐ Customer Number  
OR

☒ Registered practitioner(s) name/registration number listed below

Place Customer Number  
Bar Code Label here

Name	Registration Number	Name	Registration Number
JOANNE M. GIESSER	32,838		
JACK L. TRIBBLE	32,633		

Direct all correspondence to: ☒ Customer Number or Bar Code Label

000210

Name	JOANNE M. GIESSER				
Address	Merck & Co., Inc. - Patent Department				
Address	P.O. Box 2000, RY60-30				
City	Rahway	State	NJ	ZIP	07065-0907
Country	USA	Telephone	(732)594-3046	Fax	(732)594-4720


I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. 1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Name of Sole or First Inventor:

☐ A petition has been filed for this unsigned inventor

Given Name (first and middle [if any])  
KATHRIN U.





Family Name or Surname  
JANSEN

Inventor's Signature				Date	13 December 1999		
Residence: City	DOYLESTOWN	State	PA	Country	USA	Citizenship	GERMAN
Post Office Address	Merck & Co., Inc., P.O. Box 2000						
City	Rahway	State	NJ	ZIP	07065-0907		

☒ Additional inventors are being named on the 1 supplemental Additional Inventors(s) sheet(s) PTO/SB/02A attached hereto.

## DECLARATION AND POWER OF ATTORNEY

ADDITIONAL INVENTOR(S)  
Supplemental Sheet

Name of Additional Joint Inventor, if any:		<input type="checkbox"/> A petition has been filed for this unsigned inventor					
Given Name (first and middle [if any])		Family Name or Surname					
MARK D.		YEAGER					
Inventor's Signature					Date	13 Dec 1999	
Residence: City	DOYLESTOWN	State	PA	Country	USA	Citizenship	US
Post Office Address	Merck & Co., Inc., P.O. Box 2000						
City	Rahway	NJ	State	NJ	ZIP	07065-0907	
Name of Additional Joint Inventor, if any:		<input type="checkbox"/> A petition has been filed for this unsigned inventor					
Given Name (first and middle [if any])		Family Name or Surname					
PAUL M.		KELLER					
Inventor's Signature					Date	12/13/99	
Residence: City	LANSDALE	State	PA	Country	USA	Citizenship	US
Post Office Address	Merck & Co., Inc., P.O. Box 2000						
City	Rahway	NJ	State	NJ	ZIP	07065-0907	
Name of Additional Joint Inventor, if any:		<input type="checkbox"/> A petition has been filed for this unsigned inventor					
Given Name (first and middle [if any])		Family Name or Surname					
ROBERT S.		LOWE					
Inventor's Signature					Date	13 Dec 99	
Residence: City	HARLEYSVILLE	State	PA	Country	USA	Citizenship	US
Post Office Address	Merck & Co., Inc., P.O. Box 2000						
City	Rahway	NJ	State	NJ	ZIP	07065-0907	
Name of Additional Joint Inventor, if any:		<input type="checkbox"/> A petition has been filed for this unsigned inventor					
Given Name (first and middle [if any])		Family Name or Surname					
MIGUEL		ASTE-AMEZAGA					
Inventor's Signature					Date	December 13, 1999	
Residence: City	DREXEL HILL	State	PA	Country	USA	Citizenship	PERU
Post Office Address	Merck & Co., Inc., P.O. Box 2000						
City	Rahway	NJ	State	NJ	ZIP	07065-0907	